
PACE-IIT. MEDICAL

ACE of PACE





Factorization

- Which is the greatest number amongst $2^{1/2}$, $3^{1/3}$, $8^{1/8}$ and $9^{1/9}$? 1.

- (b) $8^{1/8}$

- (d) $2^{1/2}$
- If $a = \sqrt{25 + 4\sqrt{39}}$ and $b = \sqrt{25 4\sqrt{39}}$, then find the value of $a^3 + b^3$. 2.
 - (a) $104\sqrt{13}$
- (b) $117\sqrt{13}$
- (d) $98\sqrt{13}$
- Suppose m and n are positive real numbers such that $m\sqrt{m} + n\sqrt{n} = 183$ and $m\sqrt{n} + n\sqrt{m} = 182$. 3. Find the value of 9(m + n)
 - (a) 273
- (b) 331
- (c) 365
- (d)380

Linear Equation in Two Variables

- 4 boys and 3 girls spent Rs. 120 on the average, of which boys spend Rs. 150 on the average, then 4. the average amount spent by girls is
 - (a) Rs. 80
- (b) Rs. 60
- (c) Rs. 90
- (d) Rs. 100
- The sum of the digits of a two digit number is 14. If 18 is subtracted from the number, digits are 5. reversed. Find the number.
 - (a) 86

- (b) 77
- (c) 68
- (d)76

Quadratic Equations

- If $\sqrt{11\sqrt{11\sqrt{11......\infty}}} = 14641^x$, then find x. 6.
 - (a) 11

(b) 4

(c) $\frac{1}{4}$

(d) None of these

- If $x^2 3x + 1 = 0$, then the value of $x^5 + \frac{1}{x^5}$ 7.
 - (a) 87
- (b) 123
- (c) 135
- (d) 201
- If any polynomial f(x) is divided by $x^2 9$, then remainder is 3x + 2. If it is divided by (x 3) the 8. remainder will be:
 - (a) -7
- (b) 7

- (c) 11
- (d) -11
- If $2019^x + 2019^{-x} = 3$, then the value of $\sqrt{\frac{2019^{6x} 2019^{-6x}}{2019^x 2019^{-x}}}$ is 9.
 - (a) 3

- (b) 6
- (d) 12
- If α and β are the roots of the equation $x^2 + (\sqrt{\alpha})x + \beta = 0$ then the values of α and β are $(\alpha, \beta \neq 0)$ 10.
 - (a) $\alpha = 1$, $\beta = -2$
- (b) $\alpha = 2, \beta = -2$ (c) $\alpha = 1, \beta = -1$
- (d) $\alpha = -1$, $\beta = 1$
- If $\frac{\alpha}{\beta}$ and $\frac{\beta}{\alpha}$ are the real roots of $px^2 + qx + r = 0$, then 11.
 - (a) p = r
- (b) $q^2 = pq$ (c) $q^2 = 2pq$
- (d) none of these

12.	If one root of the equation $x^2 + px + 12 = 0$ is 4, while the equation $x^2 + px + q = 0$ has equal roots, then the value of 'q' is					
	(a) 49/4	(b) 12	(c) 3	(d) 4		
13.				-3x+5 whichever exists		
	(a) The minimum val	ue is $\frac{9}{10}$	(b) The minimum va	lue is $\frac{11}{4}$		
	(c) The maximum val	lue is $\frac{9}{10}$	(d) The maximum value is $\frac{11}{4}$			
14.		naximum value of $3 - 6x$				
	(a) $\frac{17}{8}$	(b) $\frac{33}{8}$	(c) $\frac{21}{8}$	(d) none of these		
15.	Find the range of the	values of x which satisfy				
	(a) $(-2,11)$	(b) $(2,11)$	(c) $\left(-\infty,-1\right)\cup\left(2,11\right)$	(d) $(-8, -2) \bigcup [11, \infty)$		
16.	Roots of $3^{x} + 3^{-x} = \frac{10^{x}}{2}$	$\frac{1}{3}$ are				
	(a) 0.1	(b) 1, −1	(c) $0, -1$	(d) None of these		
17.	If $x = \frac{1}{x + \frac{1}{x + \frac{1}{x + \dots \infty}}}$, the $x =$					
	(a) $\pm \frac{1}{\sqrt{2}}$		(c) 0	(d) ±1		
18.	The values of x satisf	Tying the equation is $\frac{x-1}{x-2}$	$x + \frac{x-3}{x-4} = 3\frac{1}{3}$; $x \neq 2,4$			
	(a) $3, \frac{3}{2}$	(b) $5, \frac{5}{2}$	(c) $7, \frac{7}{2}$	(d) 9, $\frac{9}{2}$		
19.	If $x = \sqrt{1 + \sqrt{1 + \sqrt{1 + \dots}}}$	${\infty}$, then x is equal to				
	(a) $\frac{1+\sqrt{5}}{2}$	(b) $\frac{1-\sqrt{5}}{2}$	$(c) \frac{1 \pm \sqrt{5}}{2}$	$(d) \frac{1 \pm \sqrt{3}}{2}$		
20.	Positive value of <i>p</i> fo will be	r which equation $x^2 + px$	$+64 = 0$ and $x^2 - 8x + y$	0 = 0 will both have real roots		
	(a) p≥16	(b) $p \le 16$	(c) $p = 16$	(d) none of these		
21.	Two numbers differ b	by 4 and their product is 1 (b) 10, 14	92. The numbers are (c) 11, 15	(d) 13, 18		

(b) $\frac{5}{4}$

22.

(a) $\frac{3}{4}$

If -4 is a root of the quadratic equation $x^2 + px - 4 = 0$ and the quadratic equation $x^2 + px + k = 0$ has equal roots, the value of k is

(c) $\frac{7}{4}$

(d) $\frac{9}{4}$

22	The volume of work which	x+3
23.	The values of x of which	$\frac{1}{x^2 - 3x - 54} \ge 0$ are

(a)
$$(-6, -3] \cup (-1, 6)$$

$$(b)[-6,-3] \cup [9,\infty]$$

$$(a) \left(-6, -3\right] \bigcup \left(-1, 6\right) \quad (b) \left[-6, -3\right] \cup \left[9, \infty\right] \qquad \quad (c) \left(-6, -3\right] \cup \left(9, \infty\right) \quad (d) \left(-6, \infty\right)$$

(d)
$$(-6,\infty)$$

The values of x for which $-2x-4 \le (x+2)^2 \le -2x-1$ is satisfied are 24.

$$(a) \left[-5, -1 \right]$$

(b)
$$[-5,0]$$

(c)
$$[-5, -4] \cup [-2, +1]$$

(d)
$$[-5,-4] \cup [-2,-1]$$

If $\frac{x^2 + x - 12}{x^2 - 3x + 2} < 0$, then x lies in _____. 25.

$$(a)(-4,3)$$

(b)
$$(-4,2)$$

(c)
$$[-4,1] \cup [2,3]$$

(c)
$$[-4,1] \cup [2,3]$$
 (d) $(-4,1) \cup (2,3)$

For what value of $x : -3x^2 + 5x - 12$ has maximum value? 26.

(a)
$$-\frac{5}{3}$$
 (b) $\frac{5}{6}$

(b)
$$\frac{5}{6}$$

(c)
$$-\frac{5}{6}$$

(d)
$$\frac{5}{3}$$

If $y^2 + 6y - 3m = 0$ and $y^2 - 3y + m = 0$ have a common root, then find the possible values of m. 27.

(a)
$$0, -\frac{27}{16}$$
 (b) $0, \frac{-81}{16}$ (c) $0, \frac{81}{16}$ (d) $0, \frac{27}{16}$

(b)
$$0, \frac{-81}{16}$$

(c)
$$0, \frac{81}{16}$$

(d)
$$0, \frac{27}{16}$$

For what value of a, the sum of roots of the equation $x^2 + 2(2-a-a^2)x - a^2 = 0$ is zero 28.

(a)
$$1, 2$$

(b)
$$1, -2$$

$$(c)-1, 2$$

$$(d) -1, -2$$

The difference between the roots of the equation $x^2 - 7x - 9 = 0$ is 29.

(b)
$$\sqrt{85}$$

(d)
$$2\sqrt{85}$$

Polynomial

If $f(x) = x^4 + ax^3 + bx^2 + cx + d$ is a polynomial such that f(1) = 5, f(2) = 10, f(3) = 15, f(4) = 20. 30.

Find the value of $\frac{f(12)+f(-8)}{100}$

If $N = \sqrt[3]{4} + \sqrt[3]{2} + 1$, then the value of $\frac{1}{N^3} + \frac{3}{N^2} + \frac{3}{N}$ is 31.

(d) 1

If $\sqrt{p} - \sqrt{q} = 20$, then the maximum value of $\left(\frac{p - 5q}{100}\right)$ is: 32.

(a) 5

(b) 10

(d) 25

How many time, graph of the polynomial $f(x) = x^3 - 1$ will intersect x-axis: 33.


(a) 0

(b) 1

(c) 2

(d) 4

34.	F(x) is a polynomial in x. When $F(x)$ is divided by $(x-2)$, the remainder obtained is 3, when the same polynomial is divided by $(x-3)$, the remainder obtained is 2. What is the remainder when $F(x)$ is divided by $(x-3)$ $(x-2)$?						
	(a) $-x + 5$	(b) $-\frac{5}{3}x + 7$	(c) 0	(d) 5			
35.	Find the other zero of t (a) -3	the polynomial $x^3 + 3x^2$ (b) 3	-2x-6, if two of its zer (c) 2	Properties are $-\sqrt{2}$ and $\sqrt{2}$ (d) None of these			



37. Value of
$$x \left[\left(1 + \frac{1}{x} \right) \left(1 + \frac{1}{x+1} \right) \left(1 + \frac{1}{x+2} \right) - 1 \right]$$
 is

(a) 3 (b) $2x$ (c) $5x$ (d) 1

38. If
$$x + y + z = 1$$
, $x^2 + y^2 + z^2 = 2$ and $x^3 + y^3 + z^3 = 3$ then the value of xyz is _____.
(a) $1/5$ (b) $1/6$ (c) $1/7$ (d) $1/8$

39. If pqr = 1, then the value of
$$\left(\frac{1}{1+p+q^{-1}} + \frac{1}{1+q+r^{-1}} + \frac{1}{1+r+p^{-1}}\right)$$
(a) 0 (b) pq (c) 1 (d) pqr

40. If
$$a = \frac{4xy}{x+y}$$
, the value of $\frac{a+2x}{a-2x} + \frac{a+2y}{a-2y}$ in most simplified form is

(a) 0 (b) 1 (c) -1 (d) 2

41. If
$$x + \frac{1}{x} = 5$$
, then $x^3 - 5x^2 + x + \frac{1}{x^3} - \frac{5}{x^2} + \frac{1}{x} = \dots$
(a) -5 (b) 0 (c) 5

42. If
$$x = \frac{1}{1+\sqrt{2}}$$
, then value of $x^2 + 2x + 3$ is:
(a) 3 (b) 0 (c) 4 (d) 1

Progression

- 43. In a certain A.P., 5 times the 5^{th} term is equal to 8 times the 8^{th} term, then its 13^{th} term is (a) 0 (b) -1 (c) -12 (d) -13
- 44. If 7th and 13th term of an A.P. be 34 and 64 respectively, then its 18th term is (a) 87 (b) 88 (c) 89 (d) 90
- 45. 7th tern of an A.P. is 40, then the sum of first 13 terms is
 (a) 53 (b) 520 (c) 1040 (d) 2080
- 46. The first term of an A.P. is 2 and common difference is 4. The sum of its 40 terms will be (a) 3200 (b) 1600 (c) 200 (d) 2800

47.	The sum of the first first term is (a) 1	at and third term of an A.	P. is 12 and the product of (c) 4	of first and second term is 24, the (d) 6			
	(a) 1	(0) 0	(C) T	(u) 0			
48.	If the sum of the finn is equal to	rst 2n terms of 2, 5, 8 i	s equal to the sum of the	first n terms of 57, 59, 61, then			
	(a) 10	(b) 12	(c) 11	(d) 13			
49.	If the sum of the 10 common difference		nes to the sum of its 5 term	ms, then the ratio of first term and			
	(a) 1:2	(b) 2:1	(c) 2:3	(d) $3:2$			
50.	If the sum of first	p terms, first q terms ar	d first r terms of an A.P	be x, y and z respectively, then			
	$\frac{x}{p}(q-r)+\frac{y}{q}(r-p)$	$+\frac{z}{r}(p-q)$ is					
	(a) 0	(b) 2	(c) pqr	$(d)\frac{8xyz}{pqr}$			
51.	If $\log 2, \log(2^n - 1)$	and $\log(2^n + 3)$ are in A	P., then n =				
	(a) $\frac{5}{2}$	(b) log ₂ 5		$(d)\frac{3}{2}$			
52.	If m th terms of the m =	sequence 163 + 167 +16	9 + and 103 + 110 +	117 + 124 + be equal, then			
	(a) 11	(b) 12	(c) 13	(d) 15			
53.	If the numbers a, b	, c, d, e form an A.P., the	en the value of $a - 4b + 6$	c-4d+e is:			
	(a) 0	(b) 2a	(c) 2c	(d) 2e			
54.	The sum of first four terms of an A.P. is 56 and sum of its last four terms is 112. If its first term is 11						
	then number of its (a) 10	(b) 11	(c) 12	(d) None of these			
55.		P. is 9 and the difference of the sequence then n		cond term is 20. If the number			
	(a) equal to 499	(b) equal to 500	(c) equal to 501	(d) can have no value			
56.			1, 2, 3, If the some	positive integers m, n we have			
	$T_m = \frac{1}{n}$ and $T_n = \frac{1}{m}$						
	(a) $\frac{1}{mn}$	(b) $\frac{1}{m} + \frac{1}{n}$	(c) 1	(d) 0			
57.	If $\log_3 2, \log_3 \left(2^x - \frac{1}{2}\right)$	(-5) and $\log_3\left(2^x - \frac{7}{2}\right)$ are	in A.P. then $x =$				
	(a) 2	(b) 3	(c) 4	(d) 2 or 3			
58.	-	_		sitive common difference. If their			
	sum is 10020, then (a) 2002	find the smallest possib (b) 2004	le value of the last term. (c) 2006	(d) 2008			

59.	0	•	n arithmetic progression. could be the length of th (c) 1275	If the lengths of the sides of the shortest side? (d) 1150				
60.	terms are three in S_1 that are there in S_2 ?							
	(a) 29	(b) 30	(c) 31	(d) 32				
61.	Find the sum of all the (a) 96750	multiples of 6 between 2 (b) 95760	00 and 1100. (c) 97560	(d) 97650				
62.	The middle term of the (a) 44	progression 4, 9, 14, (b) 49	, 104 is (c) 59	(d) 54				
63.	If the 9 th term of an A.P (a) 1:2	P. be zero, then the ratio (b) 2:1	of its 29 th and 19 th term i (c) 1:3	s: (d) 3:1				
64.	If fourth term of an A.P difference is:	. is thrice its first term a	nd seventh term -2 (thir	d term) =1, then its common				
	(a) 1	(b) 2	(c) -2	(d) 3				
65.	If the ratio of sum of n (a) 16:7	terms of two A.P's is (3r (b) 7:16	n+8): (7n+15), then th (c) 7: 12	the ratio of 12 th terms is (d) 12 : 5				
66.	the product of the secon	nd and third is 2:3, then	the smallest part is	the first and the fourth is to				
	(a) 1	(b) 2	(c) 3	(d) 4				
Logic	al Reasoning							
67.	20 m, the turns towards		e then turns towards Sou	rds North. After walking ath and walks 5 m. How far is				
	(a) 10 m, East	(b) 10 m, South-East	(c) 10 m, West	(d) 10 m, North-East				
68.	Given below are two di the face with 4?	fferent orientations of a	die. What will be the nur	mber on the face opposite to				
	$\begin{bmatrix} 1 & 6 \\ 1 & 3 \end{bmatrix}$	(L) 5	(-) 2	(1) 2				
	(a) 4	(b) 5	(c) 2	(d) 3				
69.		en, P is thirteenth from the between P and R, if R is (b) 13		from the right end. How (d) 15				
70.	The average marks of 50 students in a class was found to be 64. If the marks of two students were incorrectly entered as 38 and 42 instead of 83 and 24, respectively, then what is the correct average? (a) 64.54 (b) 62.32 (c) 61.24 (d) 61.86							
71.	Which number will rep. 47, 51, 55, 63, 87, ?	lace the question mark (?) in the given series?					
	(a) 190	(b) 183	(c) 121	(d) 155				

72.	KLIMC, JJFIX,	Which letter-cluster will replace the question mark (?) in the given series? KLIMC, JJFIX, IHCES, HFZAN, ?						
	(a) HEWWI	(b) GDUVI	(c) GDWWI	(d) GDXWJ				
73.	'P % Q' means 'P \ & Q' means If 'A @ B # C \	P' is the brother of 'Q' 'P (P' is the father of 'Q'. 'P' is the mother of 'Q'. & E @ F \ % R, then which	h of the following stater	ment is correct.				
	(a) C is the broth		(b) A is the cousing					
	(c) F is the broth	ier of R.	(d) B is the mater	nal aunt of E.				
74.	'A - B' means 'A 'A × B' means 'A 'A ÷ B' means 'A	s the mother of B'. A is the brother of B'. 'A is the father of B'. A is the daughter of B'. $J \div S + R$. then which of the of S	ne following statement is (b) Y is son of S	s not correct?				
	(c) J is daughter		(d) P is paternal u	ancle of R				
	Statements All facebook are Some whatsapp No paytm is Tel Conclusions Some Telegram Some Whatapp (a) Only I follow (c) Neither I nor	e whatsapp. are Paytm. egram. is a Facebook. is not Telegram.	(b) Only II follow (d) Both I and II f					
76.	The product of the ages of Anil and Bishnu is 240. If twice the age of Bishnu is more than							
	*	yrs. What was Bishnu's age	9					
	(a) 10 year	(b) 15 year	(c) 11 year	(d) 13 year				
77.		sons P, Q, R, S and T, T is ut younger than Q. Who is (b) T		ger than T, S is older than R and P (d) Q				
70	How many rectangles are there in the following figure?							
70.	How many recta	ingles are there in the follo	wing figure?					
73. 74. 75.		7						
	(a) 8	(b) 18	(c) 17	(d) 20				
	(, -	(-) -0	\-/ -/	(-,				

Answer key:

1. (c)	2. (d)	3. (c)	4. (a)	5. (a)	6. (c)	7. (b)	8. (c)	9. (d)	10. (a)
11. (a)	12. (a)	13. (b)	14. (b)	15. (a)	16. (b)	17. (a)	18. (b)	19. (a)	20. (c)
21. (a)	22. (d)	23. (c)	24. (d)	25. (d)	26. (b)	27. (d)	28. (b)	29. (b)	30. (d)
31. (d)	32. (a)	33. (b)	34. (a)	35. (a)	36. (b)	37. (a)	38. (b)	39. (c)	40. (d)
41. (b)	42. (c)	43. (a)	44. (c)	45. (b)	46. (a)	47. (c)	48. (c)	49. (a)	50. (a)
51. (b)	52. (c)	53. (a)	54. (b)	55. (c)	56. (c)	57. (b)	58. (c)	59. (c)	60. (c)
61. (d)	62. (d)	63. (b)	64. (b)	65. (b)	66. (b)	67. (a)	68. (c)	69. (c)	70. (a)
71. (b)	72. (c)	73. (b)	74. (d)	75. (b)	76. (a)	77. (b)	78. (b)		